



UNIVERSITY
OF WOLLONGONG
AUSTRALIA

University of Wollongong
Research Online

Faculty of Engineering and Information Sciences -
Papers: Part A

Faculty of Engineering and Information Sciences

2015

Modeling secondary particle tracks generated by high-energy protons in water

Francisco Blanco

Universidad Complutense de Madrid

Antonio Munoz

CIEMAT

Diego T. Almeida

Universidade Nova De Lisboa

F. Ferreira Da Silva

Universidade Nova De Lisboa

Paulo Limao-Vieira

Universidade Nova de Lisboa

See next page for additional authors

Publication Details

Blanco, F., Munoz, A., Almeida, D. T., Da Silva, F., Limao-Vieira, P., Verkhovtsev, A., Ellis-Gibbins, L., Krupa, K., Traore, A., Garcia, G. et al (2015). Modeling secondary particle tracks generated by high-energy protons in water. *Journal of Physics: Conference Series*, 1-1.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:
research-pubs@uow.edu.au

Modeling secondary particle tracks generated by high-energy protons in water

Abstract

We present interaction probability data of low-energy secondary electrons and positrons produced due to the proton impact. The probability distribution functions serve as input data for the Low Energy Particle Track Simulation (LEPTS) approach which allows one to include the effect of low-energy species in medical applications of radiation and in ion-beam cancer therapy, in particular.

Keywords

protons, water, high, energy, generated, modeling, tracks, particle, secondary

Disciplines

Engineering | Science and Technology Studies

Publication Details

Blanco, F., Munoz, A., Almeida, D. T., Da Silva, F., Limao-Vieira, P., Verkhovtsev, A., Ellis-Gibblings, L., Krupa, K., Traore, A., Garcia, G. et al (2015). Modeling secondary particle tracks generated by high-energy protons in water. *Journal of Physics: Conference Series*, 1-1.

Authors

Francisco Blanco, Antonio Munoz, Diego T. Almeida, F. Ferreira Da Silva, Paulo Limao-Vieira, Alexey Verkhovtsev, L Ellis-Gibblings, K Krupa, A Traore, and G Garcia

Modeling secondary particle tracks generated by high-energy protons in water

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 032092

(<http://iopscience.iop.org/1742-6596/635/3/032092>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 203.10.91.92

This content was downloaded on 28/09/2015 at 00:20

Please note that [terms and conditions apply](#).

Modeling secondary particle tracks generated by high-energy protons in water

F. Blanco^a, A. Muñoz^b, D. Almeida^c, F. Ferreira da Silva^c, P. Limão-Vieira^c,
A. Verkhovtsev^{d,e,1}, L. Ellis-Gibblings^d, K. Krupa^d, A. Traore^d, and G. García^{d,f}

^a Departamento de Física Atómica, Molecular y Nuclear, Universidad Complutense de Madrid, 28040 Madrid, Spain

^b Centro de Investigaciones, Energéticas Medioambientales y Tecnológicas (CIEMAT), Avenida Complutense 22, 28040 Madrid, Spain

^c Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal

^d CSIC, Instituto de Física Fundamental, Serrano 113-bis, 28006 Madrid, Spain

^e MBN Research Center, Altenhöferallee 3, 60438 Frankfurt am Main, Germany

^f Centre for Medical Radiation Physics, University of Wollongong, Wollongong, NSW 2522, Australia

Synopsis We present interaction probability data of low-energy secondary electrons and positrons produced due to the proton impact. The probability distribution functions serve as input data for the Low Energy Particle Track Simulation (LEPTS) approach which allows one to include the effect of low-energy species in medical applications of radiation and in ion-beam cancer therapy, in particular.

The Low Energy Particle Track Simulation (LEPTS) code [1] is a powerful complementary tool to include the effect of low-energy electrons and positrons in medical applications of radiation. In particular, for ion-beam cancer treatments it provides a detailed description of the role of the secondary electrons abundantly generated in the vicinity of the Bragg peak, as well as the possibility of using transmuted positron emitters (¹¹C, ¹⁵O) as a complement for ion-beam dosimetry.

In this study, we present interaction probability data, derived from experimental and theoretical studies, of low-energy secondary species produced due to the proton impact. The probability distribution functions, which serve as input data for the model, are also complemented with a comprehensive review of experimental [2] and theoretical [3, 4] cross sections available in the literature. We will also update the corrective factors for solid and liquid environments calculated by means of the IAM-SCAR method [5, 6]. Using these data, single electron and positron tracks in liquid water and pyrimidine have been simulated to provide information about energy deposition as well as the number and type of interactions taking place in any selected "nanovolume" of the irradiated area.

supported by the Spanish Ministry of Economía y Productividad (Project FIS2012-31320). We also acknowledge partial funding from the Portuguese Foundation for Science and Technology (FCT-MCTES), research grant PEst-OE/FIS/UI0068/2011, the EU/ESF COST Actions Nano-IBCT - MP1002 and CELINA CM-1301, and from the FP7 Multi-ITN Project "Advanced Radiotherapy, Generated by Exploiting Nanoprocesses and Technologies" (ARGENT) (Grant Agreement n°608163).

References

- [1] F. Blanco *et al* 2013 *Eur. Phys. J. D* **67** 199
- [2] M. E. Rudd, Y.-K. Kim, T. Märk, J. Schou, N. Stolterfoht, and L.H. Toburen, *Secondary Electron Spectra from Charged Particle Interactions* (International Commission on Radiation Units and Measurements, Bethesda, MD, 1996) (ICRU 55)
- [3] P. de Vera, R. Garcia-Molina, I. Abril, and A. V. Solov'yov 2013 *Phys. Rev. Lett.* **110** 148104
- [4] P. de Vera, R. Garcia-Molina, and I. Abril 2015 *Phys. Rev. Lett.* **114** 018101
- [5] F. Blanco and G. García 2003 *Phys. Rev. A* **67** 022701
- [6] F. Blanco and G. García, 2004 *Phys. Lett. A* **330** 230

Acknowledgments: This work has been

¹E-mail: verkhovtsev@iff.csic.es

